

Non-standard Confidence Sets for Ratios and Tipping Points with Applications to Dynamic Panel Data

Jean-Thomas Bernard ^{*} Ba Chu[†] Lynda Khalaf [‡]

Marcel Voia[§]

Preliminary & Incomplete: Do Not Cite or Circulate

Abstract

We study estimation uncertainty when the object of interest contains one or more ratios of parameters. The ratio of parameters is a discontinuous parameter transformation; it has been shown that traditional confidence intervals often fail to cover this true ratio with very high probability. Constructing confidence sets for ratios using Fieller's method is a viable solution as the method can avoid the discontinuity problem. This paper proposes an extension of the multivariate Fieller method beyond standard estimators, focusing on asymptotically mixed normal estimators that commonly arise in dynamic panel polynomial regression with persistent covariates. We discuss the cases where the underlying estimators converge to various distributions, depending on the persistence level of the covariates. We show that the asymptotic distribution of the pivotal statistic used for constructing a Fieller's confidence set remains a standard Chi-squared distribution regardless of rates of convergence, thus the rates are being 'self-normalized' and can be unknown. A simulation study illustrates the finite sample properties of the proposed method in a dynamic polynomial panel. Our method is demonstrated to work well in small samples, even when the persistence coefficient is unity.

^{*} Department of Economics, University of Ottawa. Mailing address: 2292 Edwin Crescent, Ottawa, Ontario K2C 1H7, Canada. TEL: 613-562-5800 ext. 1374; FAX: 613-562-5999; e-mail: jbernard3@uottawa.ca .

[†] Department of Economics, Carleton University, Carleton University. Mailing address: Loeb Building 1125 Colonel By Drive, Ottawa, Ontario, K1S 5B6 Canada. Tel (613) 520-2600-3546; FAX: (613)-520-3906. email: Ba.Chu@carleton.ca.

[‡] Department of Economics, Carleton University, and Centre de Recherche de l'Environnement, de l'Agroalimentaire, des Transports et de l'Énergie (CREATE), Centre interuniversitaire de recherche en économie quantitative (CIREQ). Mailing address: Loeb Building 1125 Colonel By Drive, Ottawa, Ontario, K1S 5B6 Canada. Tel (613) 520-2600-8697; FAX: (613)-520-3906. email: Lynda.Khalaf@carleton.ca.

[§] Department of Economics, Carleton University and Centre for Monetary and Financial Economics, Carleton University. Mailing address: Loeb Building 1125 Colonel By Drive, Ottawa, Ontario, K1S 5B6 Canada. Tel (613) 520-2600-3546; FAX: (613)-520-3906. email: Marcel.Voia@carleton.ca.

1 Introduction

Estimating or testing parameter ratios is an important issue in statistics and econometrics. From a theoretical perspective, inference problems arising from non-linearity with such transformations have attracted a great deal of interest; for references in statistics, see, for example, Zerbe, Laska, Meisner and Kushner (1982), Read (1983), Buonaccorsi (2001), Frantz (2007), and Ulrike and Franz (2009). From an empirical perspective, and more specifically in economics, ratios are parameters of interest in various applications involving elasticities or tipping points, for example with familiar "U" or inverted "U" shaped curves: Kuznet, Laffer, Rahn, Engel, Beveridge curves, as well as statistical Phillips, Yield and wage curves. In this paper, we focus on parameter ratios that are obtained from dynamic Panel data models.

In general, there are two basic approaches to estimating and assessing ratios. The first one employs a Wald-type approach, and is known as the "Delta" method [as explained in Appendix D]. This method suits asymptotically normal panel data estimators, provided of course underlying regularity conditions prevail. However, it is becoming increasingly clear from the literature that Wald-type methods raise identification problems.¹ Even when a ratio's numerator and denominator are well identified, the ratio is not well defined when its denominator approaches zero. Consequently, the distribution of standard test statistics becomes irregular, so usual tests and confidence intervals are incorrectly sized, or (said differently) usual asymptotic standard errors understate sampling uncertainty. So even if standard errors estimated using usual methods are narrow, they still provide a spurious assessment of the true uncertainty. In fact, Bolduc, Khalaf and Yelou (2010) document coverage rates collapsing to zero, that is, estimated intervals missing the unknown true value in all Monte Carlo replications, for empirically relevant scenarios.

The second approach – which may be traced back to Fieller (1954) – avoids this problem, at least in principle, using a pivotal statistic that is robust to identification as an alternative to

¹On identification problems, their consequences and possible corrections, see Dufour (2003), Staiger and Stock (1997), Wang and Zivot (1998), Zivot, Startz and Nelson (1998), Dufour and Jasiak (2001), Kleibergen (2002, 2005), Stock, Wright and Yogo (2002), Moreira (2003), Dufour and Taamouti (2005, 2007), Andrews, Moreira and Stock (2006), Antoine and Lavergne (2012).

a Wald-type one that requires identification. To the best of our knowledge, applications of the Fieller method with Panel data are scarce: Bernard, Gavin, Khalaf and Voia (2015) discussed an empirical application of the environmental Kuznet curve. Furthermore, a formal analysis of the method with persistent data is unavailable even in univariate contexts. Bernard, Idoudi, Khalaf and Yelou (2007) are a notable exception, as Monte Carlo evidence supporting the Fieller method is provided in a univariate dynamic regression, even close to the unit root boundary. In the absence of supportive theory, this result motivates further work. In time series there is work that deals with such discontinuities: Phillips (2014), Mikusheva (2007, 2012), Gorodnichenko, Mikusheva & Ng (2012). We thus revisit dynamic contexts including panel data, which as is well known, pose more serious challenges than univariate regressions. In particular for dynamic panels we extend the work of Pesaran, Shin and Smith (1999) and consider polynomial panels that span a wide range of applications; from persistence to discontinuous limiting distributions (e.g. unit root or the far-from-unity case).

As the stationarity property of polynomial regressors is often not modeled, or checked adequately the analysis of polynomial panels is interesting in its own right. We propose a parsimonious set of assumptions that preserves the stability restriction of long run equations as in Pesaran, Shin and Smith (1999) and prove that the MLE estimators converge to mixed normality at different rates. We effectively extend the multivariate Fieller method beyond standard estimators; and in the context of dynamic polynomial panels, we show that the asymptotic distribution of Fieller's statistic still remains a standard Chi-squared distribution, regardless of the convergence rates of estimates, thus the rates are being 'self-normalized' and can be unknown.

Finally, we conduct an extensive simulation study, driving persistence parameters close to boundaries, with various choices for N and T using a design based on a well know empirical example, the case of an environmental Kuznet curve. Results reveal that the delta method cannot be salvaged in dynamic Panels. The Fieller method works well with GMM methods when persistence is controlled and N is large. Fieller's method based on our likelihood based method works very well, even with unit roots, and interestingly, even when N is large relative to T .

This paper is structured as follows. Section 2 presents a general Fieller’s theorem for asymptotically mixed-normal estimators. Section 3 studies the problem of constructing Fieller’s confidence set for ratios of the parameters characterizing the long-run relationship in an error-correction representation of a dynamic heterogeneous polynomial panel. Asymptotic theory is derived for the case of fixed N and large T . Section 4 summarizes our simulation findings, and Section 5 concludes the paper. Proofs of main theorems and lemmas as well as other materials of technical flavour can be found in four appendices at the end of this paper.

1.1 Notation

The following notation is used in the paper: X denotes a scalar, \mathbf{X} is used to represent a vector or a matrix and C_0 is a generic constant that may vary from one context to another. For two random sequences, say a_T and b_T , one often writes $a_T \ll b_T$ a.s. if and only if $P(\lim_{T \uparrow \infty} |a_T/b_T| = \text{const.}) = 1$, and $a_T \ll b_T$ w.p. if and only if $\lim_{T \uparrow \infty} P(|a_T/b_T| < \xi) = 1$, where ξ can be some almost-sure bounded random variable; $o_p(\cdot)$ and $O_p(\cdot)$ are standard symbols for stochastic orders of magnitude. \xrightarrow{p} denotes convergence in probability and \xrightarrow{d} denotes convergence in distribution. $\|\cdot\|$ denotes the Euclidean norm of matrices and $\lambda_1(\mathbf{X})$ represents the minimum eigenvalue of a square matrix, \mathbf{X} . \mathbf{I}_n stands for the identity matrix of size n . $\lfloor x \rfloor$ denotes the largest integer less than or equal to x .

2 Mixed-Normality based Fieller’s Theorem

Consider a parametric model with parameters of interest defined by a vector, $\boldsymbol{\theta} = (\theta_1, \dots, \theta_p)^\top$. Let $\boldsymbol{\theta}_0 \in \Theta \subset \mathbb{R}^p$, where Θ is a compact parameter space, represent the true parameters; and for a given data sample of size T , one can estimate $\boldsymbol{\theta}_0$ by $\hat{\boldsymbol{\theta}} = (\hat{\theta}_1, \dots, \hat{\theta}_p)^\top$. We first make some assumptions about the asymptotic distribution of $\hat{\boldsymbol{\theta}}$. (Note that Assumptions 2.1 and 2.2 below are independent of each other, so are the notations.)

Assumption 2.1. $\hat{\boldsymbol{\theta}}$ is asymptotically normal as $T \uparrow \infty$, in the sense that $\mathbf{D}_T(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \xrightarrow{d}$

$N(0, \Sigma_0(\boldsymbol{\theta}_0))$ uniformly over Θ , where \mathbf{D}_T is a diagonal matrix consisting of normalizing factors that diverge to infinity with T and may differ from one another; and $N(0, \Sigma_0(\boldsymbol{\theta}_0))$ represents a tight family of Gaussian random variables with the asymptotic variance-covariance matrix, $\Sigma_0(\boldsymbol{\theta}_0)$, which is the probability limit of a matrix of normalized sample statistics, $\mathbf{D}_T^{-1} \widehat{\Sigma}_T \mathbf{D}_T^{-1}$.

Assumption 2.2. $\widehat{\boldsymbol{\theta}}$ is asymptotically mixed normal as $T \uparrow \infty$ such that

- (a) $\mathbf{D}_T(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \xrightarrow{d} \widetilde{\Sigma}_0^{-1/2}(\boldsymbol{\theta}_0)N(0, \mathbf{I}_p)$ uniformly over Θ , where \mathbf{D}_T is a diagonal matrix consisting of normalizing factors that diverge to infinity with T and may differ from one another; and $\widetilde{\Sigma}_0^{-1/2}(\boldsymbol{\theta}_0)N(0, \mathbf{I}_p)$ represents a tight family of Gaussian random variables with $\widetilde{\Sigma}_0^{-1} \equiv \widetilde{\Sigma}_0^{-1}(\boldsymbol{\theta}_0)$ being some random asymptotic variance-covariance matrix that is independent of $N(0, \mathbf{I}_p)$;
- (b) $\widetilde{\Sigma}_0$ is the probability limit of a matrix of normalized sample statistics, $\mathbf{D}_T^{-1} \widehat{\Sigma}_T \mathbf{D}_T^{-1}$, such that $\mathbf{D}_T^{-1} \widehat{\Sigma}_T \mathbf{D}_T^{-1} \xrightarrow{p} \widetilde{\Sigma}_0$.

Our objects of interest are the ratios $\boldsymbol{\rho} = (\rho_1, \dots, \rho_q)^\top$ with $\rho_i = \frac{\mathbf{L}_i^\top \boldsymbol{\theta}}{\mathbf{K}^\top \boldsymbol{\theta}}$ for $i = 1, \dots, q \leq p - 1$, where $\mathbf{L}_1, \dots, \mathbf{L}_q$, and \mathbf{K} are $q + 1$ nonstochastic and linearly independent $p \times 1$ vectors.

Theorem 1. Let either Assumption 2.1 or 2.2 hold. Then the $1 - \alpha$ asymptotic uniform simultaneous confidence sets, $CS_T(\boldsymbol{\rho}; \alpha)$, for $\boldsymbol{\rho}_0$, defined via the inverse relationship $\lim_{T \uparrow \infty} \inf_{\boldsymbol{\theta}_0 \in \Theta} P_{\boldsymbol{\theta}_0}(\boldsymbol{\rho} \in CS_T(\boldsymbol{\rho}; \alpha)) \geq 1 - \alpha$, can be obtained by inverting the following Wald-type test statistic for the null hypothesis $H_0: (\mathbf{L}^\top - \boldsymbol{\rho}_0 \mathbf{K}^\top) \boldsymbol{\theta}_0 = 0$, where $\mathbf{L} = (\mathbf{L}_1, \dots, \mathbf{L}_q)$ and $\mathbf{K}_\rho = \mathbf{K} \boldsymbol{\rho}_0^\top$.

$$\mathcal{W}(\boldsymbol{\rho}_0) = \widehat{\boldsymbol{\theta}}^\top (\mathbf{L} - \mathbf{K} \boldsymbol{\rho}_0^\top) \left((\mathbf{L} - \mathbf{K} \boldsymbol{\rho}_0^\top)^\top \widehat{\Sigma}_T^{-1} (\mathbf{L} - \mathbf{K} \boldsymbol{\rho}_0^\top) \right)^{-1} (\mathbf{L} - \mathbf{K} \boldsymbol{\rho}_0^\top)^\top \widehat{\boldsymbol{\theta}}$$

If the distributional convergence is not uniform in either Assumption 2.1 or 2.2, then one can only construct the $1 - \alpha$ asymptotic pointwise simultaneous confidence sets, $CS_T(\boldsymbol{\rho}; \alpha)$, for $\boldsymbol{\rho}_0$, defined via the inverse relationship $\lim_{T \uparrow \infty} P_{\boldsymbol{\theta}_0}(\boldsymbol{\rho} \in CS_T(\boldsymbol{\rho}; \alpha)) \geq 1 - \alpha$ for every $\boldsymbol{\theta}_0 \in \Theta$.

Therefore, a closed-form expression for $CS(\boldsymbol{\rho}; \alpha)$ can be derived by utilizing the same argument as in Section 4 of Bolduc, Khalaf and Yelou (2010).

Proof. First, an application of the uniform continuous mapping theorem yields that

$$(\mathbf{L}^\top - \mathbf{K}_\rho^\top) \mathbf{D}_T \boldsymbol{\theta} \xrightarrow{d} \left((\mathbf{L} - \mathbf{K}_\rho)^\top \tilde{\boldsymbol{\Sigma}}_0^{-1} (\mathbf{L} - \mathbf{K}_\rho) \right)^{1/2} N(0, \mathbf{I}_p) \text{ under } H_0.$$

By replacing the unknown $\tilde{\boldsymbol{\Sigma}}_0$ with $\mathbf{D}_T^{-1} \boldsymbol{\Sigma}_T \mathbf{D}_T^{-1}$, an application of Lemma 3 in Ogasawara and Takahishi (1951) yields the Wald-type statistic

$$\mathcal{W}(\boldsymbol{\rho}_0) = \hat{\boldsymbol{\theta}}^\top \mathbf{D}_T (\mathbf{L} - \mathbf{K}_\rho) \left((\mathbf{L} - \mathbf{K}_\rho)^\top \left(\mathbf{D}_T^{-1} \hat{\boldsymbol{\Sigma}}_T \mathbf{D}_T^{-1} \right)^{-1} (\mathbf{L} - \mathbf{K}_\rho) \right)^{-1} (\mathbf{L} - \mathbf{K}_\rho)^\top \mathbf{D}_T \hat{\boldsymbol{\theta}} \xrightarrow{d} \chi^2(q). \quad (2.1)$$

One can then obtain the simultaneous confidence sets for $\boldsymbol{\rho}_0$ as follows:

$$CS(\boldsymbol{\rho}; \alpha) = \{ \boldsymbol{\rho} \in \mathbb{R}^q : \mathcal{W}(\boldsymbol{\rho}) \leq c_{q,\alpha} \},$$

where $c_{q,\alpha}$ is the $(1 - \alpha)$ critical value of the χ^2 distribution with q degrees of freedom. By applying some simple transformations to the null hypothesis,

$$(\mathbf{L}^\top - \boldsymbol{\rho}_0 \mathbf{K}^\top) \boldsymbol{\theta}_0 = \mathbf{0}, \quad \Leftrightarrow (\mathbf{L}^* - \mathbf{K}^* \boldsymbol{\rho}_0^\top)^\top \mathbf{D}_T \boldsymbol{\theta}_0 = \mathbf{0}$$

where $\mathbf{L}^* = \mathbf{D}_T^{-1} \mathbf{L}$ and $\mathbf{K}^* = \mathbf{D}_T^{-1} \mathbf{K}$, we obtain in view of (2.1) that

$$\begin{aligned} \mathcal{W}(\boldsymbol{\rho}_0) &= \hat{\boldsymbol{\theta}}^\top \mathbf{D}_T (\mathbf{L}^* - \mathbf{K}^* \boldsymbol{\rho}_0^\top) \left((\mathbf{L}^* - \mathbf{K}^* \boldsymbol{\rho}_0^\top)^\top \left(\mathbf{D}_T^{-1} \hat{\boldsymbol{\Sigma}}_T \mathbf{D}_T^{-1} \right)^{-1} (\mathbf{L}^* - \mathbf{K}^* \boldsymbol{\rho}_0^\top) \right)^{-1} (\mathbf{L}^* - \mathbf{K}^* \boldsymbol{\rho}_0^\top)^\top \mathbf{D}_T \hat{\boldsymbol{\theta}} \\ &\xrightarrow{d} \chi^2(q). \end{aligned}$$

The matrices, \mathbf{D}_T , of normalizing factors in the above equation cancel out so that

$$\mathcal{W}(\boldsymbol{\rho}_0) = \hat{\boldsymbol{\theta}}^\top (\mathbf{L} - \mathbf{K} \boldsymbol{\rho}_0^\top) \left((\mathbf{L} - \mathbf{K} \boldsymbol{\rho}_0^\top)^\top \hat{\boldsymbol{\Sigma}}_T^{-1} (\mathbf{L} - \mathbf{K} \boldsymbol{\rho}_0^\top) \right)^{-1} (\mathbf{L} - \mathbf{K} \boldsymbol{\rho}_0^\top)^\top \hat{\boldsymbol{\theta}}.$$

The main theorem then follows. □

3 Dynamic Panel Polynomial Error Correction Models

Suppose that we have observations of some random variables, $y_{i,t}$, $X_{i,t}$ and $\mathbf{Z}_{i,t}$, across time periods, $t = 1, \dots, T$, and individuals, $i = 1, \dots, N$. Let the observations be generated from the following error correction model:

$$\Delta y_{i,t} = \phi_i(y_{i,t-1} - \boldsymbol{\beta}^\top \mathbf{W}_{i,t}) + \sum_{j=1}^{p-1} \lambda_{i,j} \Delta y_{i,t-j} + \sum_{j=0}^{q_x-1} \gamma_{i,j} \Delta X_{i,t-j} + \sum_{j=0}^{q_z-1} \boldsymbol{\alpha}_{i,j}^\top \Delta \mathbf{Z}_{i,t-j} + \mu_i + \epsilon_{i,t}, \quad (3.1)$$

where $\mathbf{W}_{i,t} = (\mathbf{Z}_{i,t}^\top, X_{i,t}, X_{i,t}^2, \dots, X_{i,t}^{k_x})^\top$, with $\mathbf{Z}_{i,t}$ being of dimension $k_z \times 1$, represent vectors of explanatory variables; μ_i and $\epsilon_{i,t}$ are the fixed effects and the random errors respectively; $\lambda_{i,j}$, $\gamma_{i,j}$, and $\boldsymbol{\alpha}_{i,j}$ denote the coefficients of the lagged explanatory variables; and $\boldsymbol{\beta}$ represents the regression coefficients. Conditions imposed on the dynamics of the error process and of the covariates in the d.g.p. defined by (3.1) are summarized in Assumption 3.1.

Assumption 3.1. *The innovations ϵ_i are orthogonal to both \mathbf{W}_i and \mathbf{X}_i . In addition, given i , $X_{i,t}$ is $I(1)$ and can be represented as $X_{i,t} = \sum_{s=1}^t \zeta_{i,s}$ for some zero-mean innovations, $\{\zeta_{i,t}, i = 1, \dots, N, t = 1, \dots, T\}$, which are independent across the individuals and stationary, are strongly mixing across the time periods with the mixing coefficient satisfying the condition stated in Lemma 2. The same assumptions about $X_{i,t}$ are also imposed on $\mathbf{Z}_{i,t} = \sum_{s=1}^t \boldsymbol{\xi}_{i,s}$.*

In addition, Assumption 3.2 below allows (3.1) to have a long-run relationship, $y_{i,t} = \boldsymbol{\beta}^\top \mathbf{W}_{i,t} + \nu_{i,t}$, where $\nu_{i,t}$ is a stationary process.

Assumption 3.2. *The process $y_{i,t}$ has a unit root for each i , and the lag polynomial $\sum_{j=1}^{p-1} \lambda_{i,j} z^j = 1$ has roots outside the unit circle.*

We then focus on the long-run relationship between $y_{i,t}$ and $\mathbf{W}_{i,t}$ in (3.1). Let's denote by $\boldsymbol{\varphi} = (\boldsymbol{\beta}^\top, \boldsymbol{\phi}^\top, \boldsymbol{\sigma}^\top)^\top$, where $\boldsymbol{\phi} = (\phi_1, \dots, \phi_N)^\top$ and $\boldsymbol{\sigma} = (\sigma_1^2, \dots, \sigma_N^2)^\top$, the parameters of interest, which are assumed to lie in the interior of some parameter spaces. We shall assume throughout this section that all the parameter spaces are compact, and the log-likelihood maximization is carried out on these compact spaces. Let $p^* = \max(p, q_x, q_z)$, define $(T - p^*) \times 1$ vectors, $\Delta \mathbf{y}_{i,-j} =$

$(\Delta y_{i,p^*-j}, \dots, \Delta y_{i,T-j})^\top$ and $\Delta \mathbf{X}_{i,-j} = (\Delta X_{i,p^*-j}, \dots, \Delta X_{i,T-j})^\top$, a $(T-p^*) \times k_z$ matrix, $\Delta \mathbf{Z}_{i,-j} = (\Delta \mathbf{Z}_{i,p^*-j}, \dots, \Delta \mathbf{Z}_{i,T-j})^\top$, and a $(T-p^*) \times 1$ vector of random errors, $\boldsymbol{\epsilon}_i = (\epsilon_{i,p^*}, \dots, \epsilon_{i,T})^\top$. Moreover, define

$$\mathbf{U}_i = (\Delta \mathbf{y}_{i,-1}, \dots, \Delta \mathbf{y}_{i,-p+1}, \Delta \mathbf{X}_i, \dots, \Delta \mathbf{X}_{i,-q_x+1}, \Delta \mathbf{Z}_i, \dots, \Delta \mathbf{Z}_{i,-q_z+1}, \boldsymbol{\iota}_T),$$

where $\boldsymbol{\iota}_T$ is the $(T-p^*) \times 1$ unit vector, and

$$\boldsymbol{\Lambda}_i = (\lambda_{i,1}, \dots, \lambda_{i,p-1}, \gamma_{i,0}, \dots, \gamma_{i,q_x-1}, \boldsymbol{\alpha}_{i,0}^\top, \dots, \boldsymbol{\alpha}_{i,q_z-1}^\top, \mu_i)^\top.$$

Note at this point that \mathbf{U}_i is of dimension $(T-p^*) \times k_u$ with $k_u = p + q_x + q_z k_z$, and $\boldsymbol{\Lambda}_i$ is of dimension $k_u \times 1$. One can now write (3.1) into the following matrix form:

$$\mathbf{y}_i = \phi_i(\mathbf{y}_{i,-1} - \mathbf{W}_i \boldsymbol{\beta}) + \mathbf{U}_i \boldsymbol{\Lambda}_i + \boldsymbol{\epsilon}_i,$$

where $\mathbf{W}_i = (\mathbf{Z}_i, \mathbf{X}_i, \mathbf{X}_i^2, \dots, \mathbf{X}_i^{k_x})$ with $\mathbf{Z}_i = (\mathbf{Z}_{i,1}, \dots, \mathbf{Z}_{i,T-p^*})^\top$ and $\mathbf{X}_i^\ell = (X_{i,1}^\ell, \dots, X_{i,T-p^*}^\ell)^\top$ for $\ell = 1, \dots, k_x$. The log-likelihood function is given by

$$L_T(\boldsymbol{\varphi}) = -\frac{T}{2} \sum_{i=1}^N \ln(2\pi\sigma_i^2) - \frac{1}{2} \sum_{i=1}^N \frac{1}{\sigma_i^2} (\Delta \mathbf{y}_i - \phi_i \boldsymbol{\xi}_i(\boldsymbol{\beta}))^\top \mathbf{P}_i (\Delta \mathbf{y}_i - \phi_i \boldsymbol{\xi}_i(\boldsymbol{\beta})),$$

where $\boldsymbol{\xi}_i(\boldsymbol{\beta}) = \mathbf{y}_{i,-1} - \mathbf{W}_i \boldsymbol{\beta}$ and $\mathbf{P}_i = \mathbf{I}_T - \mathbf{U}_i (\mathbf{U}_i^\top \mathbf{U}_i)^{-1} \mathbf{U}_i^\top$ with \mathbf{I}_T being the $(T-p^*) \times (T-p^*)$ identity matrix. Note that by using the orthogonality of \mathbf{P}_i to \mathbf{U}_i , one obtains for the true parameter $\boldsymbol{\varphi}_0$,

$$\begin{aligned} \frac{1}{T} (L_T(\boldsymbol{\varphi}_0) - L_T(\boldsymbol{\varphi})) &= \frac{1}{2} \sum_{i=1}^N \frac{\sigma_{0,i}^2 - \sigma_i^2}{\sigma_i^2 \sigma_{0,i}^2} \left(\frac{\boldsymbol{\epsilon}_i^\top \mathbf{P}_i \boldsymbol{\epsilon}_i}{T} - \sigma_{0,i}^2 \right) + \frac{1}{2} \sum_{i=1}^N \left(\frac{\sigma_{0,i}^2}{\sigma_i^2} - \ln \left(\frac{\sigma_{0,i}^2}{\sigma_i^2} \right) - 1 \right) \\ &\quad + \frac{1}{2T} \sum_{i=1}^N \frac{1}{\sigma_i^2} \left((\Delta \mathbf{y}_i - \phi_i \boldsymbol{\xi}_i(\boldsymbol{\beta}))^\top \mathbf{P}_i (\Delta \mathbf{y}_i - \phi_i \boldsymbol{\xi}_i(\boldsymbol{\beta})) - \boldsymbol{\epsilon}_i^\top \mathbf{P}_i \boldsymbol{\epsilon}_i \right) \\ &= \frac{1}{2} (\mathcal{T}_{1,T}(\boldsymbol{\sigma}, \boldsymbol{\sigma}_0) + \mathcal{T}_{2,T}(\boldsymbol{\sigma}, \boldsymbol{\sigma}_0) + \mathcal{T}_{3,T}(\boldsymbol{\varphi})). \end{aligned} \quad (3.2)$$

Since $\Delta \mathbf{y}_i - \phi_i \boldsymbol{\xi}_i(\boldsymbol{\beta}) = \mathbf{U}_i \boldsymbol{\Lambda}_{0,i} + \boldsymbol{\epsilon}_i + \phi_i \mathbf{W}_i (\boldsymbol{\beta}_0 - \boldsymbol{\beta}) + (\phi_i - \phi_{0,i}) \boldsymbol{\xi}_i(\boldsymbol{\beta}_0)$, one can also write

$$\mathcal{T}_{3,T}(\boldsymbol{\varphi}) = (\boldsymbol{\theta} - \boldsymbol{\theta}_0)^\top \mathbf{G}_T (\boldsymbol{\theta} - \boldsymbol{\theta}_0) + 2(\boldsymbol{\theta} - \boldsymbol{\theta}_0)^\top \mathbf{F}_T,$$

$$\text{where } \boldsymbol{\theta} = (\boldsymbol{\beta}^\top, \boldsymbol{\phi}^\top)^\top \text{ and } \mathbf{G}_T = \mathbf{G}_T(\boldsymbol{\phi}, \boldsymbol{\sigma}) = \begin{pmatrix} \sum_{i=1}^N \frac{\phi_i^2}{\sigma_i^2} \frac{\mathbf{W}_i^\top \mathbf{P}_i \mathbf{W}_i}{T} & -\frac{\phi_1}{\sigma_1^2} \frac{\mathbf{W}_1^\top \mathbf{P}_1 \boldsymbol{\xi}_{0,1}}{T} & \dots & -\frac{\phi_N}{\sigma_N^2} \frac{\mathbf{W}_N^\top \mathbf{P}_N \boldsymbol{\xi}_{0,N}}{T} \\ -\frac{\phi_1}{\sigma_1^2} \frac{\mathbf{W}_1^\top \mathbf{P}_1 \boldsymbol{\xi}_{0,1}}{T} & \frac{1}{\sigma_1^2} \frac{\boldsymbol{\xi}_{0,1}^\top \mathbf{P}_1 \boldsymbol{\xi}_{0,1}}{T} & \mathbf{0} & \mathbf{0} \\ \vdots & \mathbf{0} & \ddots & \mathbf{0} \\ -\frac{\phi_N}{\sigma_N^2} \frac{\mathbf{W}_N^\top \mathbf{P}_N \boldsymbol{\xi}_{0,N}}{T} & \mathbf{0} & \mathbf{0} & \frac{1}{\sigma_N^2} \frac{\boldsymbol{\xi}_{0,N}^\top \mathbf{P}_N \boldsymbol{\xi}_{0,N}}{T} \end{pmatrix},$$

$$\text{where } \boldsymbol{\xi}_{0,i} = \boldsymbol{\xi}_i(\boldsymbol{\beta}_0) \text{ for } i = 1, \dots, N, \text{ and } \mathbf{F}_T = \begin{pmatrix} \sum_{i=1}^N \frac{\phi_i^2}{\sigma_i^2} \frac{\mathbf{W}_i^\top \mathbf{P}_i \boldsymbol{\epsilon}_i}{T} \\ -\frac{1}{\sigma_1^2} \frac{\boldsymbol{\xi}_{0,1}^\top \mathbf{P}_1 \boldsymbol{\epsilon}_1}{T} \\ \vdots \\ -\frac{1}{\sigma_N^2} \frac{\boldsymbol{\xi}_{0,N}^\top \mathbf{P}_N \boldsymbol{\epsilon}_N}{T} \end{pmatrix}. \text{ To work out the probability}$$

limits for the random matrices \mathbf{G}_T and \mathbf{F}_T , we need to state the following lemma:

Lemma 1. *Suppose that Assumptions 3.1 and 3.2 hold. Let's denote by*

$$\mathbf{D}_{ww,T} = \text{diag}(T^{1/2} \boldsymbol{\nu}_{k_z}^\top, T^{1/2}, \dots, T^{\ell/2}, \dots, T^{k_x/2}),$$

where $\boldsymbol{\nu}_{k_z}$ is the $k_z \times 1$ unit vector, the diagonal matrix of normalizing factors. Then,

$$\mathbf{D}_{ww,T}^{-1} \frac{\mathbf{W}_i^\top \mathbf{P}_i \mathbf{W}_i}{T} \mathbf{D}_{ww,T}^{-1} \xrightarrow{p} \mathbf{Q}_{ww,i}, \quad (3.3)$$

$$\mathbf{D}_{ww,T}^{-1} \frac{\mathbf{W}_i^\top \mathbf{P}_i \boldsymbol{\xi}_{0,i}}{T} \xrightarrow{p} \mathbf{Q}_{w\xi,i}, \quad (3.4)$$

$$\frac{\boldsymbol{\xi}_{0,i}^\top \mathbf{P}_i \boldsymbol{\xi}_{0,i}}{T} \xrightarrow{p} \mathbf{Q}_{\xi\xi,i}, \quad (3.5)$$

where $\mathbf{Q}_{ww,i}$, $\mathbf{Q}_{w\xi,i}$, and $\mathbf{Q}_{\xi\xi,i}$ are some random matrices.

Theorem 2 (Consistency). *Suppose that Assumptions 3.1 and 3.2 hold. Let $\mathbf{D}_{G,T} = \text{diag}(\mathbf{D}_{ww,T}, \mathbf{I}_N)$,*

where \mathbf{I}_N is the $N \times N$ identity matrix, and

$$\mathbf{Q}_G = \mathbf{Q}_G(\boldsymbol{\phi}, \boldsymbol{\sigma}) = \begin{pmatrix} \sum_{i=1}^N \frac{\phi_i^2}{\sigma_i^2} \mathbf{Q}_{ww,i} & -\frac{\phi_1}{\sigma_1^2} \mathbf{Q}_{w\xi,1} & \dots & -\frac{\phi_N}{\sigma_N^2} \mathbf{Q}_{w\xi,N} \\ -\frac{\phi_1}{\sigma_1^2} \mathbf{Q}_{w\xi,1} & \frac{1}{\sigma_1^2} \mathbf{Q}_{\xi\xi,1} & \mathbf{0} & \mathbf{0} \\ \vdots & \mathbf{0} & \ddots & \mathbf{0} \\ -\frac{\phi_N}{\sigma_N^2} \mathbf{Q}_{w\xi,N} & \mathbf{0} & \mathbf{0} & \frac{1}{\sigma_N^2} \mathbf{Q}_{\xi\xi,N} \end{pmatrix}.$$

Moreover, suppose that $E[|\epsilon_{i,t}|^{2+\delta}] < \infty$ for some $\delta > 0$, and $\inf_{\phi, \sigma} \lambda_1(\mathbf{Q}_G) > 0$ a.s. Then,

$$\mathbf{D}_{ww,T}(\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0) = o_p(1), \quad (\widehat{\boldsymbol{\phi}} - \boldsymbol{\phi}_0) = o_p(1), \quad \text{and} \quad (\widehat{\boldsymbol{\sigma}} - \boldsymbol{\sigma}_0) = o_p(1).$$

Theorem 3 (Asymptotic Mixed Normality). *Suppose that Assumptions 3.1 and 3.2 hold. Moreover, presume that $\lambda_1(\mathbf{Q}_G(\boldsymbol{\phi}_0, \boldsymbol{\sigma}_0)) > 0$ a.s. and $E[|\epsilon_{i,t}|^{2+\delta}] < \infty$ for some $\delta > 0$. Define*

$$\mathbf{D}_T = \text{diag}(T^{1/2} \mathbf{D}_{ww,T}, T^{1/2} \mathbf{I}_N) = \text{diag}\left(T \boldsymbol{\nu}_{k_z}^\top, T, \dots, T^{\frac{\ell+1}{2}}, \dots, T^{\frac{k_x+1}{2}}, T^{\frac{1}{2}} \boldsymbol{\nu}_N^\top\right).$$

Then,

$$\mathbf{D}_T(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \xrightarrow{d} MN(\mathbf{0}, \mathbf{Q}_G^{-1}(\boldsymbol{\phi}_0, \boldsymbol{\sigma}_0)).$$

Corollary 4. *Fieller's confidence sets for the ratios $\rho_i = \frac{\mathbf{L}_i^\top \boldsymbol{\theta}_0}{\mathbf{K}^\top \boldsymbol{\theta}_0}$ for $i = 1, \dots, m$, where \mathbf{L}_i and \mathbf{K} are some given column vectors, can be constructed by inverting a Wald's statistics for testing $H_0: \mathbf{L}_i^\top \boldsymbol{\theta}_0 - \rho_{0,i} \mathbf{K}^\top \boldsymbol{\theta}_0 = 0$ vs. $H_1: \mathbf{L}_i^\top \boldsymbol{\theta}_0 - \rho_{0,i} \mathbf{K}^\top \boldsymbol{\theta}_0 \neq 0$. Theorem 3 suggests that this Wald-type statistic is given by*

$$\begin{aligned} \mathcal{W}_T(\boldsymbol{\rho}) = \boldsymbol{\theta}^\top \mathbf{D}_T (\mathbf{L} - \boldsymbol{\rho}_0 \mathbf{K}_\rho) \left((\mathbf{L}^\top - \boldsymbol{\rho}_0 \mathbf{K}_\rho^\top) (\mathbf{D}_T^{-1} \mathbf{G}_T \mathbf{D}_T^{-1})^{-1} (\mathbf{L} - \boldsymbol{\rho}_0 \mathbf{K}_\rho) \right)^{-1} (\mathbf{L}^\top - \boldsymbol{\rho}_0 \mathbf{K}_\rho^\top) \mathbf{D}_T \boldsymbol{\theta} \\ \xrightarrow{d} \chi^2(m), \end{aligned}$$

where the matrices, \mathbf{D}_T , of normalizing factors in the above equation cancel out and

$$\mathbf{G}_T = \begin{pmatrix} \sum_{i=1}^N \frac{\phi_i^2}{\sigma_i^2} \mathbf{W}_i^\top \mathbf{P}_i \mathbf{W}_i & -\frac{\phi_1}{\sigma_1^2} \mathbf{W}_1^\top \mathbf{P}_1 \boldsymbol{\xi}_1 & \dots & -\frac{\phi_N}{\sigma_N^2} \mathbf{W}_N^\top \mathbf{P}_N \boldsymbol{\xi}_N \\ -\frac{\phi_1}{\sigma_1^2} \mathbf{W}_1^\top \mathbf{P}_1 \boldsymbol{\xi}_1 & \frac{1}{\sigma_1^2} \boldsymbol{\xi}_1^\top \mathbf{P}_1 \boldsymbol{\xi}_1 & \mathbf{0} & \mathbf{0} \\ \vdots & \mathbf{0} & \ddots & \mathbf{0} \\ -\frac{\phi_N}{\sigma_N^2} \mathbf{W}_N^\top \mathbf{P}_N \boldsymbol{\xi}_N & \mathbf{0} & \mathbf{0} & \frac{1}{\sigma_N^2} \boldsymbol{\xi}_N^\top \mathbf{P}_N \boldsymbol{\xi}_N \end{pmatrix} \text{ with } \boldsymbol{\xi}_i = \mathbf{y}_{i,-1} - \mathbf{W}_i \boldsymbol{\beta} \text{ for } i = 1, \dots, N;$$

$\mathbf{D}_T^{-1} \mathbf{G}_T \mathbf{D}_T^{-1}$ is the estimate of $\mathbf{Q}_G(\boldsymbol{\phi}_0, \boldsymbol{\sigma}_0)$; $\boldsymbol{\rho} = \text{diag}(\rho_1, \dots, \rho_m)$; $\mathbf{L} = (\mathbf{L}_1, \dots, \mathbf{L}_m)$ and $\mathbf{K}_\rho = \boldsymbol{\nu}_m^\top \otimes \mathbf{K}$ are $k_{wn} \times m$ matrices.

4 Monte-Carlo Results

This section contains a Monte Carlo simulation to demonstrate the finite-sample performance of the proposed method. Specifically, we calculate the size and power of the Fieller-based test involving a ratio of two estimated parameters. To be precise the Monte-Carlo design is based on the following transformed dynamic polynomial panel:

$$\Delta y_{i,t} = \phi_i(y_{i,t-1} - \boldsymbol{\theta}^\top \mathbf{W}_{i,t}) + \sum_{j=1}^{p-1} \lambda_{i,j} \Delta y_{i,t-j} + \sum_{j=0}^{q_x-1} \gamma_{i,j} \Delta X_{i,t-j} + \sum_{j=0}^{q_z-1} \alpha_{i,j}^\top \Delta \mathbf{Z}_{i,t-j} + \mu_i + \epsilon_{i,t},$$

$i = 1, \dots, N$ and $t = 1, \dots, T$, where $\mathbf{W}_{i,t} = (\mathbf{Z}_{i,t}^\top, X_{i,t}, X_{i,t}^2, \dots, X_{i,t}^{k_x})^\top$ with a $d_z \times 1$ vector, $\mathbf{Z}_{i,t}$, contains strictly exogenous covariates; $\alpha_{i,j}$, $\lambda_{i,j}$, and $\gamma_{i,j}$ all represent the coefficients of the short-run relationship; μ_i , $i = 1, \dots, N$, indicate the fixed effects that are strictly independent of $\epsilon_{i,t}$ and $\mathbf{W}_{i,t}$; ϕ_i , $i = 1, \dots, N$, are the coefficients of the speed of adjustment to the long-run relationship; the coefficients of the long-run relationship $\boldsymbol{\theta}$ are invariant across units; $y_{i,t}$, $X_{i,t}$ and $\mathbf{Z}_{i,t}$ all have unit roots [and the long-run relationship is stationary.]

We consider that data generating process (DGP) is a finite-order $ARDL(1,0)$ process as in Pesaran and Shin (1999), where the above model includes the ECM, a quadratic polynomial and no further lags and no $Z_{i,t-j}$:

$$\begin{aligned} \phi_i &= \phi \\ X_{it} - \psi X_{i,t-1} &= \rho(X_{it} - \psi X_{i,t-1}) + \eta_{it}, \end{aligned}$$

where the errors (ϵ_{it} , η_{it}) are serially correlated and are generated according to the following bivariate normal distribution:

$$\begin{pmatrix} \epsilon \\ \mu \end{pmatrix} \rightarrow N(0, \Omega)$$

with

$$\Omega = \begin{pmatrix} 1 & \omega_{12} \\ \omega_{12} & 1 \end{pmatrix}.$$

The parameters θ comprise of θ_0 (constant), θ_1 (of X_{it}), θ_2 (of X_{it}^2), and the covariance ω_{12} , were obtained from a real data exercise done by Khalaf et al. (2011), where an empirical estimation and inference of the Environmental Kuznets Curve (EKC) for carbon dioxide and sulfur were proposed. The y_{it} in our simulations were obtained using the data on annual per capita CO_2 emission and X_{it} was measuring per capita income, in 1000s of 2000 USD. The parameters of the DGP were obtained by employing a Dynamic Panel Polynomial Error Correction Model with fixed effects (we use a DFE abbreviation as in the Figures presented in the Appendix) on CO_2 data. The above example was considered in our simulations because the original data was highly persistent and θ_2 is weakly identified. In the simulations we keep $\theta_0, \theta_1, \theta_2$, and ω_{12} fixed and we play with the degree of persistence for the y_{it} and X_{it} by changing the parameters ϕ, ρ , and ψ . In Appendix D we present a detailed description on how to construct confidence sets for a ratio of two parameters using Delta and Fieller methods.

We use different levels of persistence for both y_{it} and X_{it} starting from low persistence for both y_{it} and X_{it} to non-stationarity of X_{it} and high persistence of y_{it} . The parameters considered in the simulations are listed in the Table 1 (see Appendix).

The results of the simulation study show how poorly the Delta Method works compared to the Fieller method when we test the existence of a ratio of two parameters. In particular we show that in presence of persistent outcome variables, combining the DFE method that estimates the parameters of the model with the Fieller method used to test the existence of a ratio of two parameters, outperform any other combination of estimation and testing considered in the simulation exercise. As an alternative case we consider the Arellano-Bond (AB) estimator that is widely used for fixed T dynamic panels. We report size and power of the test underlying both Fieller and delta-method for all the cases from Table 1, however for the exposition in the paper we present few relevant cases (all the other cases are available in a separate appendix). The results show that the combination of

DFE-Fieller achieves the correct level even in finite samples, while DFE-Delta fails for any sample size. Interesting, for this combination of parameters, both the combination of AB - Delta and AB -Fieller achieves the correct level for micro panels (large N, not highly persistent data), however the combination AB-Fieller is much stable for different sample sizes than AB-Delta. From Figure 1 we can conclude that the combination of DFE-Fieller outperforms all other combinations for any sample sizes.

Figure 2 completes the picture of the performance of the combination DFE-Fieller by showing how powerful this combination is when compared to any other combination. The results also show that AB - Delta is more powerful than AB - Fieller, but much less powerful than DFE-Fieller or DFE-Delta.

In all the other cases of this Monte-Carlo study, we observe a similar behaviour for both size and power (see Figures: 3,4,5,6 for example). Therefore, we find that DFE-Fieller proposed method works in all cases where data can be highly persistent [with nonstationary covariates] while the other methods such as DFE-Delta, AB-Fieller and AB-Delta do not.

5 Conclusion

When ratios of parameters are estimated and tested, it is important to obtain reliable confidence bounds especially when one deals with longitudinal and possible nonstationary data.

As theoretical contributions, we prove that the MLE estimators for persistent dynamic panel data models converge to mixed normality at different rates, we extend the multivariate Fieller method beyond standard estimators and apply it to ratios of parameters obtained in dynamic polynomial panels and we show that the asymptotic distribution of Fieller’s statistic remains a standard Chi-squared distribution regardless of the convergence rates of estimates.

A comprehensive Monte Carlo exercise suggest that highly persistent data require adequate estimation methods coupled with appropriate testing procedures. Using a long-run estimation approach holds promise - in the sense that it provides reliable estimates for curvatures with nonstationary

data. In addition, to answer the question whether data supports a plausible tipping point, statistical methods that account for a weakly identified tipping point should be preferred. Consequently, combining the appropriate estimation method with Fieller method to construct confidence sets for ratios of parameters of interest provides a powerful tool to a researcher because the constructed confidence sets remain valid with both persistent and less persistent data.

References

- Andrews, D. W. K (2000). Inconsistency of the bootstrap when a parameter is on the boundary of the parameter space, *Econometrica* 68, 399-405.
- Andrews, D. W. K., Moreira, M. J. and Stock, J. H. (2006). Optimal two-sided invariant similar tests for instrumental variables regression, *Econometrica* 74, 715–752.
- Antoine B. and P. Lavergne (2012). Conditional moment models under weak identification, *working paper*, Simon Fraser University.
- Arellano M. and S. Bond (1991). Some tests of specification for panel data: Monte Carlo evidence and an application to employment equations, *Review of Economic Studies* 58, 277-297.
- Bernard J.-T., Idoudi N., Khalaf L. and C. Yélou (2007). Finite sample inference methods for dynamic energy demand models, *Journal of Applied Econometrics* 22, 1211-1226.
- Bernard J.-T., Gavin M., Khalaf L. and Voia M.-C. (2015). The environmental kuznets curve: tipping points and uncertainty, *Environmental and Resource Economics*, 60(2),285–315.
- Baltagi B. 1995. *Econometric Analysis of Panel Data* (Chichester: Wiley).
- Blundell R. and S. Bond (1998). Initial conditions and moment restrictions in dynamic panel data models, *Journal of Econometrics* 87, 115-143.

- Bolduc D., Khalaf L. and C. Yelou (2010). Identification robust confidence set methods for inference on parameter ratios with application to discrete choice models, *Journal of Econometrics* 157, 317–327.
- Bruno G. S. F. (2005). Estimation and inference in dynamic unbalanced panel-data models with a small number of individuals, *The Stata Journal* 5, 473-500.
- Buonaccorsi, J. P. (2001). Fieller’s theorem, in A. H. El-Shaarawi
- Beaulieu M.-C., Dufour J.-M. and L. Khalaf (2012). Testing portfolio efficiency with an unobservable zero-beta rate and non-Gaussian distributions: an exact identification-robust approach, *Review of Economic Studies*, forthcoming.
- Dufour, J.-M. (1997). Some impossibility theorems in econometrics with applications to structural and dynamic models, *Econometrica* 65, 1365–1389.
- Dufour, J.-M. (2003). Identification, weak instruments and statistical inference in econometrics, *Canadian Journal of Economics* 36(4), 767–808.
- Dufour, J.-M. and Taamouti, M. (2005). Projection-based statistical inference in linear structural models with possibly weak instruments, *Econometrica* 73, 1351–1365.
- Dufour, J.-M. and Taamouti, M. (2007). Further results on projection-based inference in IV regressions with weak, collinear or missing instruments, *Journal of Econometrics* 139, 133–153.
- Fieller E. C. (1940). The biological standardization of insulin, *Journal of the Royal Statistical Society (Supplement)* 7, 1-64.
- Fieller E. C. (1954). Some problems in interval estimation, *Journal of the Royal Statistical Society B* 16, 175-185.
- Franz, V. H. (2007). Ratios: A short guide to confidence limits and proper use, (preprint available at <http://arxiv.org/abs/0710.2024>)

- Kiviet J. F. (1995). On bias, inconsistency, and efficiency of various estimators in dynamic panel data models , *Journal of Econometrics* 68: 53-78.
- Kleibergen, F. (2002). Pivotal statistics for testing structural parameters in instrumental variables regression, *Econometrica* 70, 1781–1803.
- Kleibergen, F. (2005). Testing parameters in GMM without assuming that they are identified, *Econometrica* 73, 1103–1123.
- Moreira, M. J. (2003). A conditional likelihood ratio test for structural models, *Econometrica* 71(4), 1027–1048.
- Nickell, S. J. (1981). Biases in dynamic models with fixed effects, *Econometrica* 49: 1417-1426.
- Pesaran, M. H. and Smith R. (1995). Estimating long-run relationships from dynamic heterogeneous panels, *Journal of Econometrics* 68(1), 79–113.
- Pesaran, M. H. and Shin, Y. (1999). An autoregressive distributed lag modelling approach to cointegration analysis, in *Econometrics and Economic Theory in the 20th Century: The Ragnar Frisch Centennial Symposium*, chapter 11, (ed.) S. Strom, Cambridge University Press, Cambridge.
- Pesaran, M.H., Shin, Y., Smith, R.P (1999). Pooled mean group estimation of dynamic heterogeneous panels, *Journal of the American Statistical Association* 94, 621 - 634.
- Read, C. B. (1983). Fieller’s theorem, in *Encyclopedia of Statistical Sciences* pp. 86–88 Wiley, New York.
- Staiger, D. and Stock, J. H. (1997). Instrumental variables regression with weak instruments, *Econometrica* 65(3), 557–586.
- Stock J. H. (2010). The other transformation in econometric practice: robust tools for inference, *Journal of Economic Perspectives* 24, 83-94.

Stock J. H., J. H. Wright and M. Yogo (2002). A survey of weak instruments and weak identification in generalized method of moments, *Journal of Business and Economic Statistics* 20, 518-529.

Ulrike von L. and Franz, V. H. (2009). A geometric approach to confidence sets for ratios: Fieller's theorem, generalizations and bootstrap, *Statistica Sinica* 19, 1095-1117.

Wang, J. and Zivot, E. (1998), Inference on structural parameters in instrumental variable regression with weak instruments, *Econometrica* 66, 1389–1404.

Windmeijer, F. 2005. A finite sample correction for the variance of linear efficient two-step GMM estimators, *Journal of Econometrics* 126(1), 25-51.

Wright, J. H. (2000), Confidence set for cointegrating coefficients based on stationarity tests, *Journal of Business and Economic Statistics* 18, 211-222.

Zerbe, G.O., Laska, E., Meisner, M. and Kushner, A.B. (1982). On multivariate confidence regions and simultaneous confidence limits for ratios, *Communications in Statistics, Theory and Methods* 11, 2401 - 2425.

Zivot, E., Startz, R. and Nelson, C. R. (1998). Valid confidence intervals and inference in the presence of weak instruments, *International Economic Review* 39, 1119–1144.

Appendix A Known Results

The following lemma contains an almost sure invariance principle for sums of mixing random vectors.

Lemma 2. *Let $\{\xi_n, n \geq 1\}$ be a weak sense stationary sequence of \mathbb{R}^d -valued random vectors, centered at expectations and having $(2 + \delta)$ -th moments with $0 < \delta \leq 1$, uniformly bounded by 1; and let \mathcal{F}_a^b represent the σ -field generated by the random vectors $\xi_a, \xi_{a+1}, \dots, \xi_b$. Suppose that $\{\xi_n, n \geq 1\}$ satisfies the following strong-mixing condition:*

$$|P(AB) - P(A)P(B)| \leq \alpha(n)$$

for all $n, k \geq 1$, all $A \in \mathcal{F}_1^k$, and $B \in \mathcal{F}_{k+n}^\infty$ such that $\alpha(n) = C_0 n^{-(1+\epsilon)(1+2/\delta)}$ for some $\epsilon > 0$. Write $\xi_n = (\xi_{n,1}, \dots, \xi_{n,d})$. Then the two series in $\gamma_{i,j} = E[\xi_{1,i}\xi_{1,j}] + \sum_{k \geq 2} E[\xi_{1,i}\xi_{k,j}] + \sum_{k \geq 2} E[\xi_{k,i}\xi_{1,j}]$ converge absolutely. Denote the matrix $(\gamma_{i,j}, 1 \leq i, j \leq d)$ by Γ . Then, we can redefine the sequence $\{\xi_n, n \geq 1\}$ on a new probability space together with Brownian motion $W(t)$ with covariance matrix Γ such that

$$\sum_{n \leq t} \xi_n - W(t) \ll t^{1/2-\lambda} \text{ a.s.}$$

for some $\lambda > 0$ depending on ϵ, δ , and d only.

Proof. See Theorem 4 in Kuelbs and Philipp (1980). □

Appendix B Proofs of Auxiliary Lemmas

Proof of Lemma 1. First, note that, in view of Assumptions 3.1 and 3.2, an application of Lemma 2 yields

$$\begin{aligned} \Delta \mathbf{y}_i^\top \Delta \mathbf{y}_i &= \sum_{t=1}^T \Delta y_{i,t}^2 \ll T \text{ w.p.}, \\ \mathbf{U}_i^\top \mathbf{U}_i &\ll T \boldsymbol{\nu}_{k_u} \boldsymbol{\nu}_{k_u}^\top \text{ w.p.}, \text{ where } \boldsymbol{\nu}_{k_u} \text{ is the } k_u \times 1 \text{ unit vector,} \\ \Delta \mathbf{y}_i^\top \mathbf{Z}_i &= \sum_{t=1}^T \mathbf{Z}_{i,t} \Delta y_{i,t} \approx \int_0^1 \mathbf{Z}_{i, \lfloor T\tau \rfloor} (y_{i, \lfloor T(\tau+d\tau) \rfloor} - y_{i, \lfloor T\tau \rfloor}) \ll T \boldsymbol{\nu}_{k_z} \text{ w.p.} \end{aligned}$$

as $y_{i, \lfloor T\tau \rfloor} - y_{i, \lfloor T(\tau+d\tau) \rfloor}$ can be approximated by a Brownian motion, $dW(\lfloor T\tau \rfloor) = W(\lfloor T(\tau+d\tau) \rfloor) - W(\lfloor T\tau \rfloor)$. And by the same argument, one also obtains

$$\begin{aligned}
\Delta \mathbf{X}_i^\top \mathbf{Z}_i &\ll T \boldsymbol{\nu}_{k_z} \text{ w.p.}, \\
\Delta \mathbf{Z}_i^\top \mathbf{Z}_i &\ll T \boldsymbol{\nu}_{k_z} \boldsymbol{\nu}_{k_z}^\top \text{ w.p.}, \\
\mathbf{Z}_i^\top \mathbf{i} &= \sum_{t=1}^T \mathbf{Z}_{i,t} \ll T^{3/2} \boldsymbol{\nu}_{k_z} \text{ w.p.}, \\
(\mathbf{X}_i^\ell)^\top \Delta \mathbf{y}_i &= \sum_{t=1}^T X_{i,t}^\ell \Delta y_{i,t} \approx \int_0^1 X_{i, \lfloor T\tau \rfloor}^\ell (y_{i, \lfloor T(\tau+d\tau) \rfloor} - y_{i, \lfloor T\tau \rfloor}) \ll T^{\frac{\ell+1}{2}} \text{ w.p.}, \\
(\mathbf{X}_i^\ell)^\top \boldsymbol{\nu}_T &\ll T^{\frac{\ell+2}{2}} \text{ w.p.}, \\
(\mathbf{X}_i^\ell)^\top \mathbf{U}_i &\ll \left(T^{\frac{\ell+1}{2}} \boldsymbol{\nu}_{k_u-1}^\top, T^{\frac{\ell+2}{2}} \right) \text{ w.p.}
\end{aligned}$$

Collecting all the above-derived rates of divergence, one can immediately show that

$$\mathbf{Z}_i^\top \mathbf{U}_i \ll (T \boldsymbol{\nu}_{k_z \times (k_u-1)}, T^{3/2} \boldsymbol{\nu}_{k_z}) \text{ w.p.},$$

where $\boldsymbol{\nu}_{k_z \times (k_u-1)}$ represents the $k_z \times (k_u - 1)$ unit matrix. Some matrix manipulations then yield

$$\begin{aligned}
\mathbf{D}_{ww,T}^{-1} \frac{\mathbf{W}_i^\top \mathbf{U}_i (\mathbf{U}_i^\top \mathbf{U}_i)^{-1} \mathbf{U}_i^\top \mathbf{W}_i}{T} \mathbf{D}_{ww,T}^{-1} &\xrightarrow{p} \mathbf{Q}_{ww,i}^{(1)}, \\
\mathbf{D}_{ww,T}^{-1} \frac{\mathbf{W}_i^\top \mathbf{W}_i}{T} \mathbf{D}_{ww,T}^{-1} &\xrightarrow{p} \mathbf{Q}_{ww,i}^{(2)}.
\end{aligned}$$

Hence, (3.3) immediately follows. In addition, note that

$$\begin{aligned}
\mathbf{Z}_i^\top \boldsymbol{\xi}_{0,i} &\ll \sum_{t=1}^T t^{1/2} \approx T^{3/2} \text{ w.p.}, \\
(\mathbf{X}_i^\ell)^\top \boldsymbol{\xi}_{0,i} &\ll \sum_{t=1}^T t^{\ell/2} = T^{\frac{\ell+2}{2}}, \text{ w.p.} \\
\mathbf{Z}_i^\top \boldsymbol{\xi}_{0,i} &\ll T^{3/2} \text{ w.p.}, \\
(\mathbf{X}_i^\ell)^\top \boldsymbol{\xi}_{0,i} &\ll T^{\frac{\ell+2}{2}} \text{ w.p.}
\end{aligned}$$

One can immediately show (3.4) and (3.5). □

Appendix C Proofs of Main Theorems

Proof of Theorem 2. We adopt the strategy used in Saikkonen (1995) and Pesaran, Shin and Smith (1998). First, define the open shrinking balls: $B_T(\boldsymbol{\beta}_0, \delta_\beta) = \{\boldsymbol{\beta} \in \Theta_\beta \subset \mathbb{R}^{k_w} : \|\mathbf{D}_{ww,T}(\boldsymbol{\beta} - \boldsymbol{\beta}_0)\| < \delta_\beta\}$, where $k_w = k_z + k_x$ and Θ_β is some compact parameter space of $\boldsymbol{\beta}_0$; $B(\boldsymbol{\phi}_0, \delta_\phi) = \{\boldsymbol{\phi} \in \Theta_\phi \subset \mathbb{R}^N : \|\boldsymbol{\phi} - \boldsymbol{\phi}_0\| < \delta_\phi\}$, where Θ_ϕ is some compact parameter space of $\boldsymbol{\phi}_0$; and $B(\boldsymbol{\sigma}_0, \delta_\sigma) = \{\boldsymbol{\sigma} \in \Theta_\sigma \subset \mathbb{R}^N : \|\boldsymbol{\sigma} - \boldsymbol{\sigma}_0\| < \delta_\sigma\}$, where Θ_σ is some compact parameter space of $\boldsymbol{\sigma}_0$. Let $B_T^c(\boldsymbol{\beta}_0, \delta_\beta)$, $B^c(\boldsymbol{\phi}_0, \delta_\phi)$, and $B^c(\boldsymbol{\sigma}_0, \delta_\sigma)$ be the complements of $B_T(\boldsymbol{\beta}_0, \delta_\beta)$, $B(\boldsymbol{\phi}_0, \delta_\phi)$, and $B(\boldsymbol{\sigma}_0, \delta_\sigma)$ respectively. Define $\mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma) = \left\{ \bigcup_{\{\delta_\beta, \delta_\phi: (\delta_\beta^2 + \delta_\phi^2)^{1/2} = \delta\}} B_T^c(\boldsymbol{\beta}_0, \delta_\beta) \times B^c(\boldsymbol{\phi}_0, \delta_\phi) \right\} \times B^c(\boldsymbol{\sigma}_0, \delta_\sigma)$. We need to show that

$$\lim_{T \uparrow \infty} P \left(\inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} \frac{1}{T} (L_T(\boldsymbol{\varphi}_0) - L_T(\boldsymbol{\varphi})) > 0 \right) = 1 \tag{C-1}$$

for every $\delta, \delta_\sigma > 0$. In view of (3.2), one obtains

$$\inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} \frac{1}{T} (L_T(\boldsymbol{\varphi}_0) - L_T(\boldsymbol{\varphi})) \geq \frac{1}{2} \left\{ \inf_{\boldsymbol{\sigma} \in B^c(\boldsymbol{\sigma}_0, \delta_\sigma)} \mathcal{T}_{1,T}(\boldsymbol{\sigma}, \boldsymbol{\sigma}_0) + \inf_{\boldsymbol{\sigma} \in B^c(\boldsymbol{\sigma}_0, \delta_\sigma)} \mathcal{T}_{2,T}(\boldsymbol{\sigma}, \boldsymbol{\sigma}_0) + \inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} \mathcal{T}_{3,T}(\boldsymbol{\varphi}) \right\}.$$

It can immediately be shown that $\inf_{\boldsymbol{\sigma} \in B^c(\boldsymbol{\sigma}_0, \delta_\sigma)} \mathcal{T}_{1,T}(\boldsymbol{\sigma}, \boldsymbol{\sigma}_0) = o_p(1)$ and $\inf_{\boldsymbol{\sigma} \in B^c(\boldsymbol{\sigma}_0, \delta_\sigma)} \mathcal{T}_{2,T}(\boldsymbol{\sigma}, \boldsymbol{\sigma}_0) > 0$. Furthermore,

$$\inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} \mathcal{T}_{3,T}(\boldsymbol{\varphi}) \geq \inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} (\boldsymbol{\theta} - \boldsymbol{\theta}_0)^\top \mathbf{G}_T (\boldsymbol{\theta} - \boldsymbol{\theta}_0) + 2 \inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} (\boldsymbol{\theta} - \boldsymbol{\theta}_0)^\top \mathbf{F}_T = \mathcal{T}_{3,a,T}(\boldsymbol{\varphi}) + 2\mathcal{T}_{3,b,T}(\boldsymbol{\varphi}).$$

Note that, by an elementary matrix inequality and Lemma 1,

$$\begin{aligned} \mathcal{T}_{3,a,T}(\boldsymbol{\varphi}) &= \inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} (\boldsymbol{\theta} - \boldsymbol{\theta}_0)^\top \mathbf{D}_{G,T} [\mathbf{D}_{G,T}^{-1} \mathbf{G}_T \mathbf{D}_{G,T}^{-1}] \mathbf{D}_{G,T} (\boldsymbol{\theta} - \boldsymbol{\theta}_0) \\ &\geq \inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} \|\mathbf{D}_{G,T} (\boldsymbol{\theta} - \boldsymbol{\theta}_0)\|^2 \inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} \lambda_1 (\mathbf{D}_{G,T}^{-1} \mathbf{G}_T \mathbf{D}_{G,T}^{-1}) \\ &\geq \delta^2 \inf_{\substack{\boldsymbol{\phi} \in B(\boldsymbol{\phi}_0, \delta_\phi) \\ \boldsymbol{\sigma} \in B(\boldsymbol{\sigma}_0, \delta_\sigma)}} \lambda_1(\mathbf{Q}_G) \text{ w.p.} \end{aligned}$$

It then follows that $\mathcal{T}_{3,a,T}(\boldsymbol{\varphi}) > 0$ w.p. Moreover, by Lemma 1, one has

$$\begin{aligned} \mathbf{D}_{ww,T}^{-1} \frac{\mathbf{W}_i^\top \mathbf{P}_i E[\boldsymbol{\epsilon}_i^\top \boldsymbol{\epsilon}_i] \mathbf{P}_i \mathbf{W}_i}{T} \mathbf{D}_{ww,T}^{-1} &\xrightarrow{p} \sigma_i^2 \mathbf{Q}_{ww,i}, \\ \frac{\boldsymbol{\xi}_{0,i}^\top \mathbf{P}_i E[\boldsymbol{\epsilon}_i^\top \boldsymbol{\epsilon}_i] \mathbf{P}_i \boldsymbol{\xi}_{0,i}}{T} &\xrightarrow{p} \sigma_i^2 \mathbf{Q}_{\xi\xi,i}. \end{aligned}$$

Conditioning \mathbf{F}_T on \mathbf{W}_i , \mathbf{P}_i and $\boldsymbol{\xi}_i$, an application of the multivariate CLT to the sequence $\boldsymbol{\epsilon}_i$ yields $\mathbf{D}_{G,T}^{-1} \mathbf{F}_T = O_p(T^{-1/2}) = o_p(1)$. Since, from the way $\mathcal{B}_T(\boldsymbol{\beta}_0, \delta_\beta)$ is defined, the term $\inf_{\boldsymbol{\varphi} \in \mathcal{B}_T(\boldsymbol{\varphi}, \delta, \delta_\sigma)} (\boldsymbol{\theta} - \boldsymbol{\theta}_0)^\top \mathbf{D}_{G,T}^{-1} \boldsymbol{\iota}_{k_w+N}$ is bounded either above or below by a generic constant, which can be large but does not depend on T , it immediately follows that $\mathcal{T}_{3,b,T} = o_p(1)$. Therefore, (C-1) has been verified. \square

Proof of Theorem 3. The gradient and Hessian matrices of $L_T(\boldsymbol{\varphi})$ are given by $\frac{\partial L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\theta}} = \left(\frac{\partial L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta}^\top}, \frac{\partial L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\phi}^\top} \right)^\top$

and $\frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^\top} = \begin{pmatrix} \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta} \partial \boldsymbol{\beta}^\top} & \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta} \partial \boldsymbol{\phi}} \\ \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta} \partial \boldsymbol{\phi}} & \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\phi} \partial \boldsymbol{\phi}^\top} \end{pmatrix}$, where

$$\begin{aligned} \frac{\partial L_T(\boldsymbol{\varphi})}{\partial \phi_i} &= \frac{1}{\sigma_i^2} \boldsymbol{\xi}_i(\boldsymbol{\beta})^\top \mathbf{P}_i(\Delta \mathbf{y}_i - \phi_i \boldsymbol{\xi}_i(\boldsymbol{\beta})), \\ \frac{\partial L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta}} &= - \sum_{i=1}^N \frac{\phi_i}{\sigma_i^2} \mathbf{W}_i^\top \mathbf{P}_i(\Delta \mathbf{y}_i - \phi_i \boldsymbol{\xi}_i(\boldsymbol{\beta})), \\ \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\phi_i \phi_j} &= 0 \text{ for } i \neq j, \\ \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\phi_i^2} &= - \frac{1}{\sigma_i^2} \boldsymbol{\xi}_i(\boldsymbol{\beta})^\top \mathbf{P}_i \boldsymbol{\xi}_i(\boldsymbol{\beta}), \\ \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta} \partial \boldsymbol{\beta}^\top} &= - \sum_{i=1}^N \frac{\phi_i^2}{\sigma_i^2} \mathbf{W}_i^\top \mathbf{P}_i \mathbf{W}_i, \\ \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta} \partial \phi_i} &= - \frac{1}{\sigma_i^2} \mathbf{W}_i^\top \mathbf{P}_i(\Delta \mathbf{y}_i - \phi_i \boldsymbol{\xi}_i(\boldsymbol{\beta})) + \frac{\phi_i}{\sigma_i^2} \mathbf{W}_i^\top \mathbf{P}_i \boldsymbol{\xi}_i(\boldsymbol{\beta}). \end{aligned}$$

Since $\widehat{\boldsymbol{\varphi}}$ is consistent by Theorem 2, an application of a first-order Taylor expansion of $\frac{\partial L_T(\widehat{\boldsymbol{\varphi}})}{\partial \boldsymbol{\theta}}$ about $\boldsymbol{\theta}_0$ yields

$$0 = \frac{\partial L_T(\widehat{\boldsymbol{\varphi}})}{\partial \boldsymbol{\theta}} = \frac{\partial L_T(\boldsymbol{\theta}_0, \widehat{\boldsymbol{\sigma}})}{\partial \boldsymbol{\theta}} + \frac{\partial^2 L_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^\top} (\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0),$$

where $\boldsymbol{\theta}^*$ is some point lying on the line segment $L(\boldsymbol{\theta}_0, \widehat{\boldsymbol{\theta}}) = \{s\boldsymbol{\theta}_0 + (1-s)\widehat{\boldsymbol{\theta}} : s \in (0, 1)\} \subset \Theta_\beta \times \Theta_\phi \subset \mathbb{R}^{k_{wn}}$, where $\Theta_\beta \times \Theta_\phi$ are the compact parameter spaces of $\boldsymbol{\theta}_0$ (as defined in the proof of Theorem 2), and $k_{wn} = k_w + N$. One can then obtain

$$\mathbf{D}_T(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) = - \left[\mathbf{D}_T^{-1} \frac{\partial^2 L_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^\top} \mathbf{D}_T^{-1} \right]^{-1} \mathbf{D}_T^{-1} \frac{\partial L_T(\boldsymbol{\theta}_0, \widehat{\boldsymbol{\sigma}})}{\partial \boldsymbol{\theta}}. \quad (\text{C-2})$$

For notational brevity, let $\mathcal{I}_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}}) = \mathbf{D}_T^{-1} \frac{\partial^2 L_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^\top} \mathbf{D}_T^{-1}$. First, one needs to show that

$$\lim_{T \uparrow \infty} P(\|\mathcal{I}_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}}) - \mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0)\| > \epsilon) = 0 \text{ given some arbitrarily small } \epsilon > 0. \quad (\text{C-3})$$

Note that

$$\begin{aligned}
& P(\|\mathcal{I}_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}}) - \mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0)\| > \epsilon) \\
&= P(\|\mathcal{I}_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}}) - \mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0)\| > \epsilon \mid \boldsymbol{\theta}^* \in B_T(\boldsymbol{\beta}_0, \delta_\beta) \times B(\boldsymbol{\phi}_0, \delta_\phi), \widehat{\boldsymbol{\sigma}} \in B(\boldsymbol{\sigma}_0, \delta_\sigma)) \\
&\quad P(\boldsymbol{\theta}^* \in B_T(\boldsymbol{\beta}_0, \delta_\beta) \times B(\boldsymbol{\phi}_0, \delta_\phi), \widehat{\boldsymbol{\sigma}} \in B(\boldsymbol{\sigma}_0, \delta_\sigma)) \\
&+ P(\|\mathcal{I}_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}}) - \mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0)\| > \epsilon \mid \boldsymbol{\theta}^* \in B_T^c(\boldsymbol{\beta}_0, \delta_\beta) \times B^c(\boldsymbol{\phi}_0, \delta_\phi), \widehat{\boldsymbol{\sigma}} \in B^c(\boldsymbol{\sigma}_0, \delta_\sigma)) \\
&\quad P(\boldsymbol{\theta}^* \in B_T^c(\boldsymbol{\beta}_0, \delta_\beta) \times B^c(\boldsymbol{\phi}_0, \delta_\phi), \widehat{\boldsymbol{\sigma}} \in B^c(\boldsymbol{\sigma}_0, \delta_\sigma)),
\end{aligned}$$

where the balls $B_T(\boldsymbol{\beta}_0, \delta_\beta)$, $B(\boldsymbol{\phi}_0, \delta_\phi)$, and $B(\boldsymbol{\sigma}_0, \delta_\sigma)$ are defined in the proof of Theorem 2. Since $\lim_{T \uparrow \infty} P(\boldsymbol{\theta}^* \in B_T^c(\boldsymbol{\beta}_0, \delta_\beta) \times B^c(\boldsymbol{\phi}_0, \delta_\phi), \widehat{\boldsymbol{\sigma}} \in B^c(\boldsymbol{\sigma}_0, \delta_\sigma)) = 0$ for every $\boldsymbol{\theta}^*$ lying on the line segment $L(\boldsymbol{\theta}_0, \widehat{\boldsymbol{\theta}})$ by Theorem 2, one has

$$\lim_{T \uparrow \infty} P(\|\mathcal{I}_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}}) - \mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0)\| > \epsilon) \leq \lim_{T \uparrow \infty} P\left(\sup_{\substack{\boldsymbol{\theta} \in B_T(\boldsymbol{\beta}_0, \delta_\beta) \times B(\boldsymbol{\phi}_0, \delta_\phi) \\ \boldsymbol{\sigma} \in B(\boldsymbol{\sigma}_0, \delta_\sigma)}} \|\mathcal{I}_T(\boldsymbol{\theta}, \boldsymbol{\sigma}) - \mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0)\| > \epsilon\right) \quad (\text{C-4})$$

for some arbitrarily small numbers, δ_β , δ_ϕ and δ_σ . An application of Lemma 1 and some inequalities for matrices yields

$$\begin{aligned}
\sup_{\substack{\boldsymbol{\phi} \in B(\boldsymbol{\phi}_0, \delta_\phi) \\ \boldsymbol{\sigma} \in B(\boldsymbol{\sigma}_0, \delta_\sigma)}} \left\| \mathbf{D}_{\gamma, k_w}^{-1} \left(\frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta} \partial \boldsymbol{\beta}^\top} - \frac{\partial^2 L_T(\boldsymbol{\varphi}_0)}{\partial \boldsymbol{\beta} \partial \boldsymbol{\beta}^\top} \right) \mathbf{D}_{\gamma, k_w}^{-1} \right\| &\leq C_0 (\delta_\phi^2 + \delta_\sigma^2)^{\frac{1}{2}} \sum_{i=1}^N \|\mathbf{Q}_{ww, i}\|, \\
\sup_{\substack{\boldsymbol{\phi} \in B(\boldsymbol{\phi}_0, \delta_\phi) \\ \boldsymbol{\sigma} \in B(\boldsymbol{\sigma}_0, \delta_\sigma)}} \left\| \mathbf{D}_{\gamma, k_w}^{-1} \left(\frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\beta} \partial \boldsymbol{\phi}^\top} - \frac{\partial^2 L_T(\boldsymbol{\varphi}_0)}{\partial \boldsymbol{\beta} \partial \boldsymbol{\phi}^\top} \right) \mathbf{I}_N T^{1/2} \right\| &\leq C_0 \left(\delta_\beta \sum_{i=1}^N \|\mathbf{Q}_{ww, i}\| + (\delta_\phi^2 + \delta_\sigma^2)^{\frac{1}{2}} \sum_{i=1}^N \|\mathbf{Q}_{w\xi, i}\| \right), \\
\sup_{\substack{\boldsymbol{\phi} \in B(\boldsymbol{\phi}_0, \delta_\phi) \\ \boldsymbol{\sigma} \in B(\boldsymbol{\sigma}_0, \delta_\sigma)}} \left\| \frac{\partial^2 L_T(\boldsymbol{\varphi})}{\partial \boldsymbol{\phi} \partial \boldsymbol{\phi}^\top} - \frac{\partial^2 L_T(\boldsymbol{\varphi}_0)}{\partial \boldsymbol{\phi} \partial \boldsymbol{\phi}^\top} \right\| &\leq \delta_\sigma \left(\sum_{i=1}^N \|\mathbf{Q}_{\xi\xi, i}\| \right)^{1/2},
\end{aligned}$$

where C_0 is some finite generic constant that may differ from a line to another one. An application

of the matrix inequality: $\| \begin{smallmatrix} A & C \\ C & D \end{smallmatrix} \| \leq \|A\|_2 + \sqrt{2}\|C\|_2 + \|D\|_2$ yields

$$\begin{aligned} & \sup_{\substack{\boldsymbol{\theta} \in B_T(\boldsymbol{\beta}_0, \delta_\beta) \times B(\boldsymbol{\phi}_0, \delta_\phi) \\ \boldsymbol{\sigma} \in B(\boldsymbol{\sigma}_0, \delta_\sigma)}} \|\mathcal{I}_T(\boldsymbol{\theta}, \boldsymbol{\sigma}) - \mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0)\| \\ & \leq C_0 \left((\delta_\beta + (\delta_\phi^2 + \delta_\sigma^2)^{1/2}) \sum_{i=1}^N \|\mathbf{Q}_{ww,i}\| + (\delta_\phi^2 + \delta_\sigma^2)^{1/2} \sum_{i=1}^N \|\mathbf{Q}_{w\xi,i}\| + \delta_\sigma \sum_{i=1}^N \|\mathbf{Q}_{\xi\xi,i}\| \right). \end{aligned} \quad (\text{C-5})$$

The consistency of $\widehat{\boldsymbol{\theta}}$ allows one to make δ_β , δ_ϕ , and δ_σ in (C-5) arbitrarily small such that its RHS becomes less than ϵ . In view of (C-4), (C-3) has been proved. Therefore, $\|\mathcal{I}_T(\boldsymbol{\theta}^*, \widehat{\boldsymbol{\sigma}}) - \mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0)\| = o_p(1)$. By the same argument, one can also show that

$$\left\| \mathbf{D}_T^{-1} \left(\frac{\partial L_T(\boldsymbol{\theta}_0, \widehat{\boldsymbol{\sigma}})}{\partial \boldsymbol{\theta}} - \frac{\partial L_T(\boldsymbol{\varphi}_0)}{\partial \boldsymbol{\theta}} \right) \right\| = o_p(1).$$

Moreover, by Lemma 1, one has

$$\mathcal{I}_T(\boldsymbol{\theta}_0, \boldsymbol{\sigma}_0) \xrightarrow{p} \mathbf{Q}_G(\boldsymbol{\phi}_0, \boldsymbol{\sigma}_0),$$

where \mathbf{Q}_G is given in Theorem 2. Now, notice that for each

$$E_\epsilon \left[\mathbf{D}_T^{-1} \frac{\partial L_T(\boldsymbol{\varphi}_0)}{\partial \boldsymbol{\theta}} \frac{\partial L_T(\boldsymbol{\varphi}_0)}{\partial \boldsymbol{\theta}^\top} \mathbf{D}_T^{-1} \right] = \mathbf{Q}_G(\boldsymbol{\phi}_0, \boldsymbol{\sigma}_0),$$

where the expectation is taken with respect to the joint probability density of $\boldsymbol{\epsilon}_i$. Therefore, conditioning $\mathbf{D}_T^{-1} \frac{\partial L_T(\boldsymbol{\varphi}_0)}{\partial \boldsymbol{\theta}}$ on \mathbf{W}_i , \mathbf{P}_i , and $\boldsymbol{\xi}_i(\boldsymbol{\beta}_0)$, an application of the multivariate CLT to the sequence $\boldsymbol{\epsilon}_i$ yields

$$\mathbf{D}_T^{-1} \frac{\partial L_T(\boldsymbol{\varphi}_0)}{\partial \boldsymbol{\theta}} \xrightarrow{d} N(\mathbf{0}, \mathbf{Q}_G(\boldsymbol{\phi}_0, \boldsymbol{\sigma}_0)).$$

The main theorem then follows from (C-2) and some marginal integration. □

Appendix D Confidence Set for Ratios of two Parameters

[Tipping Points]

Consider the general model $(\mathcal{Y}, \{P_\theta : \theta \in \Theta\})$, $\Theta \subset \mathbb{R}^p$, $p \geq 1$, where \mathcal{Y} is the sample space and P_θ is a probability distribution over \mathcal{Y} indexed by $\theta = (\theta_1, \theta_2, \dots, \theta_p)'$. Our object of interest are functions of θ of the form $h(\theta) = \mathbf{L}'\theta/\mathbf{K}'\theta$ where \mathbf{L} and \mathbf{K} are nonstochastic $p \times 1$ vectors. Given a sample of size T , assume a consistent and asymptotically normal estimator of θ is available $\hat{\theta} = (\hat{\theta}_1, \hat{\theta}_2, \dots, \hat{\theta}_p)' \stackrel{asy}{\approx} N(\theta, \Sigma_\theta)$ where Σ_θ is estimated consistently by $\hat{\Sigma}_\theta$. The discontinuity set $\{\theta \in \Theta : \mathbf{K}'\theta = 0\}$ is clearly non-empty. In this context, the *delta* method exploits the following regular asymptotic result:

$$h(\hat{\theta}) \stackrel{asy}{\approx} N \left(h(\theta), \frac{\partial h(\hat{\theta})}{\partial \theta'} \hat{\Sigma}_\theta \frac{\partial h'(\hat{\theta})}{\partial \theta} \right). \quad (\text{D-1})$$

For the same problem, Fieller's method inverts a Wald-type test associated with the hypothesis $\mathbf{L}'\theta - \rho_0 \mathbf{K}'\theta = 0$ for a collection of fixed ρ_0 values. For the ratio case presented in Section 2, Fieller's method involves assembling all ρ_0 values such that $\theta_1 - \rho_0 \theta_2 = 0$ is not rejected at the $\alpha\%$ using the *t*-statistic $(\hat{\theta}_1 - \rho_0 \hat{\theta}_2) / (\rho^2 \hat{v}_2 - 2\rho_0 \hat{v}_{12} + \hat{v}_1)^{1/2}$ which is asymptotically standard normal under the null hypothesis. The confidence set is thus defined as solution to following inequality in ρ_0

$$FCS(\rho; \alpha) = \left\{ \rho_0 : \left(\hat{\theta}_1 - \rho_0 \hat{\theta}_2 \right)^2 \leq z_{\alpha/2}^2 (\hat{v}_1 + \rho_0^2 \hat{v}_2 - 2\rho_0 \hat{v}_{12}) \right\}. \quad (\text{D-2})$$

This requires solving the following second-degree-polynomial inequality for ρ_0 :

$$A\rho_0^2 + 2B\rho_0 + C \leq 0 \quad (\text{D-3})$$

$$A = \hat{\theta}_2^2 - z_{\alpha/2}^2 \hat{v}_2, \quad B = -\hat{\theta}_1 \hat{\theta}_2 + z_{\alpha/2}^2 \hat{v}_{12}, \quad C = \hat{\theta}_1^2 - z_{\alpha/2}^2 \hat{v}_1. \quad (\text{D-4})$$

for real solutions ρ_0 . Except for a set of measure zero, $A \neq 0$. Similarly, except for a set of measure zero, $\Delta = B^2 - AC \neq 0$. Real roots

$$\delta_{01} = \frac{-B - \sqrt{\Delta}}{A}, \quad \delta_{02} = \frac{-B + \sqrt{\Delta}}{A}$$

exist if and only if $\Delta > 0$, so

$$FCS(\rho; \alpha) = \begin{cases} [\rho_{01}, \delta_{02}] & \text{if } A > 0 \\]-\infty, \delta_{01}] \cup [\delta_{02}, +\infty[& \text{if } A < 0 \end{cases}. \quad (\text{D-5})$$

Bolduc, Khalaf and Yelou (2010) further show that: (i) if $\Delta < 0$, then $A < 0$ and $FCS(\rho; \alpha) = R$; (ii) $FCS(\rho; \alpha)$ contains the point estimate $\hat{\rho} = \hat{\theta}_1/\hat{\theta}_2$ and thus cannot be empty, and (iii) asymptotically, Fieller's solution and the *delta* method give similar results when the former leads to an interval, *i.e.* when the denominator is far from zero.

Table 1: Parameters - for Monte Carlo Simulation

Parameters	1	2	3	4	5	6	7	8
ρ	0.2	0.8	0.9	0.99	0.2	0.8	0.9	0.99
ψ	1	1	1	1	0.2	0.2	0.2	0.2
ϕ	1	1	1	1	1	1	1	1
ω_{12}	0.85	0.85	0.85	0.85	0.85	0.85	0.85	0.85
θ_0	-0.679	-0.679	-0.679	-0.679	-0.679	-0.679	-0.679	-0.679
θ_1	0.619	0.619	0.619	0.619	0.619	0.619	0.619	0.619
θ_2	-0.007	-0.007	-0.007	-0.007	-0.007	-0.007	-0.007	-0.007

Table 2: Size and Power for $\rho = 0.2$, $\phi = 1$, $\psi = 1$

	N	Size				Power			
		T=5	T=10	T=50	T=100	T=5	T=10	T=50	T=100
PMG Delta	10	0.117	0.514	1	1	0.566	0.87	1	1
	50	0.22	0.668	1	1	0.721	0.971	1	1
	100	0.335	0.745	1	1	0.787	0.998	1	1
	150	0.406	0.792	1	1	0.832	1	1	1
PMG Fieller	10	0.066	0.061	0.055	0.057	0.781	1	1	1
	50	0.057	0.047	0.055	0.051	1	1	1	1
	100	0.05	0.056	0.067	0.063	1	1	1	1
	150	0.045	0.055	0.066	0.051	1	1	1	1
AB Delta	10	0.162	0.086	0.125	0.165	0.381	0.388	0.501	0.531
	50	0.03	0.054	0.134	0.218	0.305	0.333	0.477	0.515
	100	0.038	0.057	0.132	0.198	0.296	0.348	0.346	0.517
	150	0.053	0.05	0.121	0.119	0.322	0.356	0.308	0.265
AB Fieller	10	0.147	0.066	0.086	0.1	0.138	0.092	0.067	0.069
	50	0.053	0.078	0.054	0.097	0.032	0.071	0.055	0.064
	100	0.031	0.068	0.074	0.07	0.037	0.048	0.067	0.061
	150	0.019	0.054	0.137	0.176	0.027	0.039	0.117	0.069

Notes:

Table 3: Size and Power for $\rho = 0.9$, $\phi = 1$, $\psi = 1$

	N	Size				Power			
		T=5	T=10	T=50	T=100	T=5	T=10	T=50	T=100
PMG Delta	10	0.987	1	1	1	0.984	0.997	1	1
	50	1	1	1	1	1	0.971	1	1
	100	1	1	1	1	0.787	0.998	1	1
	150	1	1	1	1	0.832	1	1	1
PMG Fieller	10	0.068	0.067	0.053	0.049	0.781	1	1	1
	50	0.052	0.048	0.046	0.05	1	1	1	1
	100	0.062	0.056	0.065	0.05	1	1	1	1
	150	0.044	0.055	0.059	0.057	1	1	1	1
AB Delta	10	0.801	0.919	0.989	0.992	0.984	0.997	1	1
	50	0.947	1	0.953	0.998	1	1	0.999	1
	100	0.941	1	1	0.957	1	1	1	0.997
	150	0.963	1	1	1	1	1	1	1
AB Fieller	10	0.174	0.146	0.414	0.477	0.916	0.968	0.996	0.996
	50	0.06	0.225	0.237	0.673	1	1	0.954	1
	100	0.056	0.212	0.989	0.306	0.999	1	1	0.887
	150	0.049	0.215	0.998	0.999	0.999	1	1	1

Notes:

Table 4: Size and Power for $\rho = 0.2$, $\phi = 0.2$, $\psi = 1$

	N	Size				Power			
		T=5	T=10	T=50	T=100	T=5	T=10	T=50	T=100
PMG Delta	10	0.161	0.508	1	1	0.657	0.873	1	1
	50	0.383	0.814	1	1	0.863	0.999	1	1
	100	0.516	0.902	1	1	0.948	1	1	1
	150	0.581	0.94	1	1	0.974	1	1	1
PMG Fieller	10	0.064	0.066	0.056	0.058	0.956	1	1	1
	50	0.074	0.048	0.055	0.05	1	1	1	1
	100	0.057	0.059	0.044	0.063	1	1	1	1
	150	0.059	0.069	0.060	0.061	1	1	1	1
AB Delta	10	0.15	0.084	0.146	0.185	0.366	0.416	0.503	0.518
	50	0.04	0.066	0.147	0.216	0.3	0.342	0.497	0.513
	100	0.035	0.061	0.104	0.189	0.291	0.334	0.244	0.497
	150	0.037	0.048	0.108	0.08	0.285	0.333	0.165	0.162
AB Fieller	10	0.142	0.073	0.061	0.087	0.133	0.087	0.059	0.066
	50	0.05	0.072	0.175	0.098	0.05	0.105	0.056	0.061
	100	0.025	0.053	0.137	0.186	0.039	0.083	0.107	0.048
	150	0.021	0.054	0.305	0.306	0.031	0.087	0.259	0.108

Notes:

Table 5: Size and Power for $\rho = 0.9$, $\phi = 0.2$, $\psi = 1$

	N	Size				Power			
		T=5	T=10	T=50	T=100	T=5	T=10	T=50	T=100
PMG Delta	10	0.998	1	1	1	1	1	1	1
	50	1	1	1	1	1	1	1	1
	100	1	1	1	1	1	1	1	1
	150	1	1	1	1	1	1	1	1
PMG Fieller	10	0.07	0.072	0.056	0.048	1	1	1	1
	50	0.083	0.054	0.046	0.054	1	1	1	1
	100	0.092	0.078	0.05	0.056	1	1	1	1
	150	0.09	0.081	0.063	0.06	1	1	1	1
AB Delta	10	0.895	0.918	0.991	0.997	0.995	0.996	1	1
	50	0.996	1	0.956	1	1	1	0.994	1
	100	0.994	1	1	0.964	1	1	1	0.997
	150	0.997	1	1	1	1	1	1	1
AB Fieller	10	0.191	0.159	0.423	0.498	0.978	0.96	0.997	0.99
	50	0.143	0.513	0.225	0.663	1	1	0.945	1
	100	0.137	0.523	1	0.345	1	1	1	1
	150	0.116	0.57	1	1	1	1	1	1

Notes:

Table 6: Size and Power for $\rho = 0.2$, $\phi = 1$, $\psi = 0.8$

	N	Size				Power			
		T=5	T=10	T=50	T=100	T=5	T=10	T=50	T=100
PMG Delta	10	0.056	0.216	0.594	0.653	0.446	0.689	0.968	0.997
	50	0.098	0.306	0.782	0.887	0.625	0.751	1	1
	100	0.161	0.369	0.739	0.968	0.663	0.845	1	1
	150	0.194	0.44	0.772	0.916	0.702	0.868	1	1
PMG Fieller	10	0.068	0.056	0.048	0.064	0.545	1	1	1
	50	0.061	0.047	0.055	0.078	0.986	1	1	1
	100	0.06	0.061	0.06	0.102	1	1	1	1
	150	0.046	0.061	0.048	0.102	1	1	1	1
AB Delta	10	0.194	0.155	0.064	0.071	0.347	0.345	0.306	0.315
	50	0.028	0.038	0.0083	0.057	0.252	0.249	0.275	0.296
	100	0.038	0.026	0.032	0.09	0.224	0.246	0.283	0.311
	150	0.031	0.037	0.035	0.043	0.262	0.256	0.286	0.282
AB Fieller	10	0.146	0.074	0.067	0.053	0.157	0.091	0.074	0.072
	50	0.042	0.052	0.051	0.056	0.043	0.071	0.048	0.059
	100	0.039	0.046	0.019	0.345	0.037	0.046	0.07	0.048
	150	0.019	0.054	0.046	0.028	0.03	0.055	0.106	0.076

Notes:

Table 7: Size and Power for $\rho = 0.9$, $\phi = 0.2$, $\psi = 1$

	N	Size				Power			
		T=5	T=10	T=50	T=100	T=5	T=10	T=50	T=100
PMG Delta	10	0.998	1	1	1	1	1	1	1
	50	1	1	1	1	1	1	1	1
	100	1	1	1	1	18	1	1	1
	150	1	1	1	1	1	1	1	1
PMG Fieller	10	0.07	0.072	0.056	0.048	1	1	1	1
	50	0.083	0.054	0.046	0.054	1	1	1	1
	100	0.092	0.078	0.05	0.056	1	1	1	1
	150	0.09	0.081	0.063	0.06	1	1	1	1
AB Delta	10	0.895	0.918	0.991	0.997	0.995	0.996	1	1
	50	0.996	1	0.956	1	1	1	0.994	1
	100	0.994	1	1	0.964	1	1	1	0.997
	150	0.997	1	1	1	1	1	1	1
AB Fieller	10	0.191	0.159	0.423	0.498	0.978	0.96	0.997	0.99
	50	0.143	0.513	0.225	0.663	1	1	0.945	1
	100	0.137	0.523	1	0.345	1	1	1	1
	150	0.116	0.57	1	1	1	1	1	1

Notes:

Table 8: Size and Power for $\rho = 0.2$, $\phi = 0.2$, $\psi = 0.99$, $\beta = 0.5$

	N	Size				Power			
		T=5	T=10	T=50	T=100	T=5	T=10	T=50	T=100
PMG Delta	10	1	1	1	1	1	1	1	1
	50	1	1	1	1	1	1	1	1
	100	1	1	1	1	1	1	1	1
PMG Fieller	10	0.75	0.045	0.058		1	1	1	1
	50	0.045	0.047	0.047		1	1	1	1
	100	0.064	0.058	0.054		1	1	1	1
AB Delta	10	1	1	1	1	1	1	1	1
	50	1	1	1	1	1	1	1	1
	100	1	1	1	1	1	1	1	1
AB Fieller	10	0.157	0.079	0.064		1	1	1	1
	50	0.048	0.06	0.253		1	1	1	1
	100	0.027	0.049	0.091		1	1	1	1

Notes: